



Experimental investigation and thermodynamic modeling of the C-Me-O systems (where Me = Hf, Zr)

Florian Rejasse, Olivier Rapaud, Gilles Trolliard, Alexandre Maitre

► To cite this version:

Florian Rejasse, Olivier Rapaud, Gilles Trolliard, Alexandre Maitre. Experimental investigation and thermodynamic modeling of the C-Me-O systems (where Me = Hf, Zr). 15th International IUPAC Conference on High Temperature Materials Chemistry (HTMC XV), Mar 2016, Orléans, France. hal-01297096

HAL Id: hal-01297096

<https://hal.science/hal-01297096>

Submitted on 2 Apr 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Experimental investigation and thermodynamic modeling of the C-Me-O systems (where Me = Hf, Zr)

F. Réjasse¹, O. Rapaud¹, G. Trolliard¹, A. Maître¹


¹ Univ.Limoges, CNRS, ENSCI, SPCTS, UMR 7315, F-87000 Limoges France

Introduction

- Zirconium and hafnium carbide (ZrC_x and HfC_x) are considered as Ultra High Temperature Ceramics (UHTC) which can be easily oxidized under flowing dioxygen.
- Oxygen atoms can substitute carbon atoms in the crystalline lattice of carbides involving the formation of oxycarbide phases ZrC_xO_y and HfC_xO_y .
- The thermodynamic modeling of the C-Me-O ternary phase diagrams requires thermodynamic and diagrammatic data especially to determine the solubility of the oxygen into the carbide sublattice.

Experimental



Reaction mechanisms involved during the carbothermal reduction of ZrO_2 and HfO_2 ^{a, b}



Structural approach by **XRD** and **TEM** → understanding of the reaction mechanisms involved for **different advancements** ξ .

Destabilization of ZrO_2 and HfO_2

Destabilization of C




Experimental determination of solid solution boundaries by carbothermal reduction



- The amount of initial reactants is adjusted to produce several oxycarbide stoichiometries.
- Heat treatment in a graphite furnace under flowing argon at **1650°C, 1750°C and 1850°C** during 8 hours ($\xi = 1$).

→ For ZrC_xO_{1-x}




Stoichiometries in C and O of each sample computed by elemental analysis :

- Increase of the maximum solubility of O with the increase of the temperature**

- No modification of the maximum solubility of C with the temperature**

An **abacus** can be created correlating the lattice parameter and the stoichiometry of ZrC_xO_{1-x}

Temperatures	Low boundary	High boundary
1650°C	$ZrC_{0.83}O_{0.17}$	$ZrC_{0.94}O_{0.03}$
1750°C	$ZrC_{0.80}O_{0.20}$	$ZrC_{0.96}O_{0.04}$
1850°C	$ZrC_{0.78}O_{0.22}$	$ZrC_{0.94}O_{0.03}$





→ For HfC_xO_{1-x}


- Between 1650°C and 1750 °C : no evolution of the solubility of O
- At the same temperature, the stability domain of HfC_xO_{1-x} solid solution is narrower than ZrC_xO_y**

Nucleation of the MeC_xO_{1-x} phase

Co-condensation reaction of $MeO_{(g)}$ and $CO_{(g)}$ within carbon-rich areas



Evolution of the MeC_xO_{1-x} phase




The increase of the carbon amount in the crystalline structure implies the increase of the MeC_xO_{1-x} lattice parameter.

Purely reconstructive reaction → Impossible to obtain nanosized carbides by this synthesis way


Thermodynamic Modeling

Binary diagram Zr-C ^c




- FCC_A1 : $(Zr)_1(C,Va)_1 / L^0, L^1$
- C Graphite
- BCC_A2 : $(Zr)_1(C,Va)_3 / L^0$
- HCP_A3 : $(Zr)_1(C,Va)_{0.5} / L^0$
- Liquid : $(Zr,C)_1 / L^0, L^1, L^2$

Extrapolation in the ternary system and result of optimisation




Binary diagram Zr-O ^d



- FCC_C1 : $(Zr)_1(O,Va)_2(O,Va)_1 / L^0, L^1, L^2$
- BCC_A2 : $(Zr)_1(O,Va)_3 / L^0$
- HCP_A3 : $(Zr)_1(O,Va)_{0.5} / L^0$
- Liquid : $(O,ZrO_2, Zr)_1 / L^0, L^1, L^2$


At 1750°C

Binary diagram Hf-C ^e




- FCC_B1 : $(Hf)_1(C,Va)_1 / L^0, L^1$
- C Graphite
- BCC_A2 : $(Hf)_1(C,Va)_3 / L^0$
- HCP_A3 : $(Hf)_1(C,Va)_{0.5} / L^0, L^1, L^2$
- Liquid : $(Hf,C)_1 / L^0, L^1, L^2, L^3$

Extrapolation in the ternary system and result of optimisation



Binary diagram Hf-O ^f



- BCC_A2 : $(Hf)_1(O,Va)_3 / L^0$
- HCP_A3 : $(Hf)_1(O,Va)_{0.5} / L^0, L^1$
- Liquid : $(Hf,O)_1 / L^0, L^1, L^2$

At 1750°C

Summary

- The carbothermal reaction is a **purely reconstructive reaction** which operates through solid-gas equilibria.
- This reaction allowed to determine the stability domains of MeC_xO_y solid solutions. On the Zr-C-O system, the extent of the domain increases with the temperature. At the same temperature, the domain of HfC_xO_y is narrower than ZrC_xO_y .
- These **experimental data** were used to assess the **two ternary phase diagrams** and to describe the evolution of the solid solutions with the temperature